

Abundance and community structure of Mentawai primates in the Peleonan Forest, North Siberut, Indonesia

MATTHIAS WALTERT, CHRISTOPHE ABEGG, THOMAS ZIEGLER, SUSILO HADI,
DODI PRIATA and KEITH HODGES

Abstract The island of Siberut in the Mentawai Archipelago, west of Sumatra, Indonesia, supports four of the five primate species endemic to the Archipelago, all categorized as threatened on the IUCN Red List. As a baseline for future research on the ecology and conservation of Siberut's primates we used a line-transect approach to survey primates in the Peleonan forest, site of the Siberut Conservation Project in Northern Siberut. In 104 km of surveys we made a total of 391 observations of primates and estimated density and population sizes for the project's 10.7 km² main study site. Overall primate biomass density was estimated to be 697 kg km⁻² and abundances, in individuals km⁻², were: pig-tailed snub-nosed monkey *Simias concolor*, 53.1; Siberut macaque *Macaca siberu*, 16.2; Kloss's gibbon *Hylobates klossii*, 8.9; Mentawai langur *Presbytis potenziani*, 8.2. These results confirm those from earlier studies that in undisturbed and unhunted primary forests *S. concolor* is more successful than the sympatrically occurring *P. potenziani*. Results for all species suggest considerable population sizes on the island and underline its importance for the conservation of Mentawai primates.

Keywords *Hylobates klossii*, Indonesia, *Macaca siberu*, *Presbytis potenziani*, primate density, Siberut, *Simias concolor*.

Introduction

Having been isolated from other land masses for at least 500,000 years (Verstappen, 1973), the 7,800 km² Mentawai Island Archipelago west of Sumatra, Indonesia, supports numerous endemic mammals. Endemism of non-volant mammals is exceptionally high (WWF, 1982), including five endemic primate species in four genera: the monotypic pig-tailed snub-nosed langur or *simakobu*

Simias concolor, *Hylobates klossii*, the Mentawai langur *Presbytis potenziani*, and two species of macaque (*Macaca pagensis* and *Macaca siberu*). *H. klossii* occurs throughout the archipelago (Whittaker, 2005a). *M. pagensis* occurs on the three smaller islands of Sipora, North and South Pagai, and *M. siberu* is confined to the largest and most northerly island, Siberut (Kitchener & Groves, 2002; Roos *et al.*, 2003). Subspecies have been described for both langur species based on morphological characters only (Chasen & Kloss, 1927).

This high level of primate endemism is a priority for conservation (Chivers, 1986; MacKinnon, 1986; Mittermeier *et al.*, 2007) but the Mentawai primates are under threat from habitat loss resulting from extensive deforestation, both legal and illegal, for agricultural use. This has reduced habitat for primates on the Mentawai islands by at least 50% in the last 25 years (from an estimated 4,200 km² to only 2,400 km² (Chivers, 1986; Whittaker, 2005b). Furthermore, hunting primates for food is an important part of Mentawai culture. According to local hunters (pers. comm.) *S. concolor* is the easiest to hunt and its meat is the best tasting; it is therefore the main prey species (Mitchell & Tilson, 1986). The species is listed globally as one of the 25 most endangered primates (Mittermeier *et al.*, 2007).

Although all Mentawai primate taxa are categorized as threatened on the IUCN Red List (Endangered: *S. concolor*; Vulnerable: *H. klossii*, *P. potenziani*; Critically Endangered: *M. pagensis* and *M. siberu*; IUCN, 2007), there is relatively little reliable and up-to-date information on their population sizes and distribution within the archipelago (Tenaza, 1987; Paciulli, 2004). Data for Siberut, accounting for over half (4,480 km²) of the total land mass (Whitten, 1982b; Schefold, 1988) are particularly limited and no transect surveys of primate densities have been carried out. Siberut still has considerable forest cover and, in contrast to the southern islands of the archipelago, a comparatively low human population density, especially in the north (Tenaza, 1987; Fuentes & Ray, 1995; Yanuar *et al.*, 1998).

Previous estimates of the density of Siberut's primates were made by Tilson (1977), Watanabe (1981) and Whittaker (2005b). Tilson (1977) estimated densities of *S. concolor* at one site in Central Siberut, and Watanabe (1981) studied *S. concolor* and *P. potenziani* at two sites, in north and south Siberut. Whether these studies are representative is difficult to ascertain because density estimates were

MATTHIAS WALTERT (Corresponding author) Department of Conservation Biology, Centre for Nature Conservation, Georg-August-University, Von-Siebold-Strasse 2, 37075 Göttingen, Germany. E-mail mwalter@gwdg.de

CHRISTOPHE ABEGG, THOMAS ZIEGLER, SUSILO HADI and KEITH HODGES Department of Reproductive Biology, German Primate Centre, Kellnerweg 4, 37077 Göttingen, Germany.

DODI PRIATA Institute of Research and Community Empowerment, Bogor Agricultural University, Bogor, Indonesia.

Received 12 October 2006. Revision requested 30 November 2006.

Accepted 25 June 2007.

mainly inferred from home ranges within comparatively small areas of *c.* 2 km². In addition, the widely differing results from the two locations of Watanabe's study (7 and 220 individuals per km², for *S. concolor*) complicate interpretation of the data. Call monitoring was used by Whittaker (2005a, b) to calculate estimates of population size and density of *H. klossii* in northern Siberut.

Here we present data from the Peleonan forest, a 40 km² conservation area within a belt of production forest in northern Siberut, close to ongoing logging activities in the east and to the Siberut National Park in the south and west. The Peleonan forest is one of the few remaining relatively undisturbed primary rainforests on the island and is currently assigned for the research and conservation activities of the Siberut Conservation Project under an agreement with the local clans. In addition to conducting primatological research, one of the main aims of the Project is the establishment of a database to support the development of conservation measures for the region and for other areas situated within production forests. As a part of this we surveyed the four primate taxa present, and here we present data on abundance, density and biomass using line transect data collected in 2005. This data provides a baseline for future island-wide population estimates.

Study area

The *c.* 40 km² Peleonan forest is in the north-east of the island and connected to the large forest block of Siberut National Park by a wide corridor of mixed primary and secondary forest. It borders the coast to the north, extends from sea level to 180 m, and consists of primary dipterocarp rainforest dominating high ridges and hills. On slopes and lower elevations primary mixed forest, characterized mainly by Myristicaceae, Euphorbiaceae, Dilleniaceae and Dipterocarpaceae, dominates. Near the northern coast the Peleonan forest turns into waterlogged peat swamp forest, characterized by tree species of Myristicaceae and Sapotaceae, and fringed by permanently wet freshwater swamp forest dominated by *Terminalia phellocarpa* associated with feather palms, rattans, pandans and aroids (WWF, 1982). These swampy areas are separated from the sea by a narrow belt of coastal vegetation. The Siberut Conservation Project field station is located in the south of the Peleonan forest (Fig. 1) and its main study site is a circular area of 10.7 km² around the station that has been made accessible by 13 systematically placed radially oriented transects with a mean length of 1.5 km (range 1–3 km, 19.85 km in total).

Methods

Surveys for primates were carried out between 9 July and 29 December 2005 in a total survey effort of 104.1 km. Surveys were conducted over 6.30–11.30 and 15.30–18.00. Protocols,




FIG. 1 Location of Siberut (a) off West Sumatra (b), and the Siberut Conservation Project's field station and the main study site within the Peleonan forest (c).

including searching behaviour and data analysis (see below), followed those of earlier primate surveys (Waltert *et al.*, 2002) and general recommendations for line transect surveys (Peres, 1999; Buckland *et al.*, 2001; Plumptre & Cox, 2006). For each observation the species, group size and perpendicular distance from the group or subgroup's centre to the transect (measured using a laser range finder), were noted. We assessed detection probabilities by fitting half-normal detection functions with hermite adjustments to perpendicular distance data, using the software *Distance v. 4.1* (Thomas *et al.*, 2001), and calculated density and abundance for the survey site.

As radially oriented transects could potentially lead to biased density estimates by over-covering the area surrounding the station we tested whether survey parameters differed between distances from the station. For each species we calculated encounter rates, detection probability and cluster size for three transect sections, i.e. 0–500, 500–1,000 and 1,000–1,500 m from the starting points. Only *S. concolor* had a significant drop in encounter rate within the first transect section compared to the second and third (Kruskall-Wallis test $P < 0.001$). Detection probability and cluster size did not differ significantly (z -test, $P > 0.1$). To avoid a biased density estimate for this species we stratified the data by dividing the 10.7 km² study area into the inner

belt surrounding the station (a circular area with 600 m radius), and the remaining (c. 1,200 m wide) outer belt.

For all species the size-bias regression method (natural log of cluster size against estimated detection probability) was used to correct average cluster size (sub-group size noted in the field) for distance effects. Because of the dense habitat it was often not possible to measure perpendicular distances accurately beyond 30 m, and only estimation of distances was possible. We therefore compared results from data truncated to <30 m with only slightly truncated data and found that density estimates derived from slightly truncated data (75–99 m truncation distance) were not significantly different but coefficients of variation were much lower (12–28% instead of 15–40%) in all species.

The daily activity patterns of gibbons are usually reflected in encounter rates, which differ between morning and afternoon surveys. Because of the pooling robustness of the detection function such differences do not pose a problem for detection probability estimation, and hence should not lead to biased density estimates if stratified analyses are not made. We nevertheless also modelled detection probability using survey time (morning vs afternoon) as a covariate. Visual examinations of distance distributions showed there were differences in the shape of the detection functions between morning and afternoon surveys, suggesting that more cryptic behaviour in the afternoon could have led to reduced detection probability near the transect centre and that our density estimates of *H. klossii* may be conservative.

Biomass density was calculated according to the average weights of Mentawai primates (Rowe, 1996): *S. concolor*, 7.9 kg; *P. potenziani*, 6.5 kg; *M. siberu*, 9.5 kg; *H. klossii*, 5.8 kg.

Results

Average group sizes were 2.1–3.2 individuals (Table 1) but many observed *M. siberu* groups were sub-groups of larger social units. A total of 391 primate groups were encountered: *S. concolor*, 187; *M. siberu*, 71; *P. potenziani*, 67. *H. klossii*, 66. Detection functions fitted to data truncated to widths <100 m resulted in 139, 41, 36 and 34 observations for each species, respectively (Table 1). *S. concolor* was the most abundant primate species, with an average density of

TABLE 2 Mean density estimate, with 95% confidence interval (CI) and coefficient of variation of density (CV), biomass density, and population estimates (based on 95% confidence intervals of mean density), of the four primate species in the Peleonan forest, the site of the Siberut Conservation Project in Northern Siberut (Fig. 1).

Species	Mean density, km ⁻² (95% CI, % CV)	Biomass density (kg km ⁻²)	Population size, range
<i>S. concolor</i>	53.1 (38.8–72.5, 15.9)*	419	420–780
<i>P. potenziani</i>	8.2 (4.8–14.1, 26.8)	53	50–150
<i>M. siberu</i>	16.2 (9.4–27.9, 27.5)	154	100–300
<i>H. klossii</i>	8.9 (5.7–14.0, 22.4)	52	60–150

*Estimate corrected for density decrease with increasing distance from field station.

53.1 individuals km⁻² (and a biomass density of 419 kg km⁻²) and an estimated population of 570 (range 420–780; Table 2). Stratifying data by distance from the station showed that average density of *S. concolor* decreases from c. 70 km⁻² within 600 m of the station to c. 50 km⁻² elsewhere; other than this decrease there was no evidence for further falls in encounter rate (and hence density) at distances >600 m from the field station. All other primates were considerably less abundant than *S. concolor* (Table 2). Mean primate biomass was 679 kg km⁻² and, based on 95% confidence limits of density, 460–1011 kg km⁻².

Discussion

This study provides the first systematic line transect survey of primates on Siberut and the first to encompass all of its four endemic primate taxa. Given that the Peleonan forest has so far been relatively undisturbed by logging and hunting, the study provides an insight into the presumed natural structure of the endemic primate community. The most abundant primate of Siberut's forests appears to be *S. concolor*, and this species is probably capable of reaching high densities in areas with low disturbance and little hunting pressure (Watanabe, 1981). Our calculated densities for this species are approximately double those obtained by Tenaza & Fuentes (1995) on North and South Pagai, underlining the importance of the Peleonan forest for this

TABLE 1 Mean group encounter rate, size and density, and detection probability, with 95% confidence intervals (CI), for groups within truncation distance *w*, and number of groups encountered (*n*), of the four primate species in the Peleonan forest, the site of the Siberut Conservation Project in Northern Siberut (Fig. 1).

Species	Mean group encounter rate, km ⁻¹ (95% CI)	Mean group size (95%CI)	Mean group density, km ⁻² (95% CI)	Detection probability (95% CI)	<i>w</i>	<i>n</i>
<i>Simias concolor</i>	1.72 (1.46–2.01)	2.6 (2.3–2.9)	16.2 (12.5–21.1)	0.58 (0.50–0.67)	75	139
<i>Presbytis potenziani</i>	0.35 (0.22–0.55)	2.1 (1.8–2.5)	3.2 (1.9–5.4)	0.55 (0.41–0.72)	99	36
<i>Macaca siberu</i>	0.39 (0.26–0.60)	3.2 (2.4–4.1)	3.8 (2.3–6.2)	0.57 (0.43–0.74)	91	41
<i>Hylobates klossii</i>	0.33 (0.24–0.45)	2.2 (1.8–2.7)	3.5 (2.3–5.3)	0.50 (0.38–0.65)	93	34

species. Thus, conservation efforts in undisturbed areas, even in small areas such as the Peleonan forest, could be successful in securing viable populations of this species.

In contrast, the density of the other Siberut colobine, *P. potenziani*, appears to be naturally much lower, probably <50% of the density of *S. concolor*, confirming estimates made by Watanabe (1981). The reason for this large difference between the two species is unclear but could be due to their different ecologies (Tilson, 1977; Watanabe, 1981).

Whittaker (2005b) found *H. klossii* at densities >25 km⁻² in the Peleonan forest but our estimates are >50% lower. The reason for this discrepancy is difficult to determine because of the different sampling techniques used. While undercounting of gibbons can occur with line transects, even on the transect line (Brockelman & Ali, 1987), and our estimate may therefore be conservative, our interpretation is that Whittaker may have overestimated density due to a combination of low sample size, lack of systematic sampling, and underestimation of gibbon call distance. Our estimate of 8–9 *H. klossii* km⁻² is similar to that of Whitten (1982a) who estimated 10.5 individuals km⁻² based on home ranges, and to data for other *Hylobates* spp. (density ranges of 4–12 individuals km⁻²; McConkey & Chivers, 2004; Nijman, 2004), although Brockelman *et al.* (1998) reported much higher densities for *Hylobates lar* (5 groups km⁻²). Whittaker's (2005b) estimate of average group size for *H. klossii* in the Peleonan forest (3.7) is also unusually high compared to other estimates for *H. klossii* (Whitten, 1982a; Paciulli, 2004) and to the other four locations surveyed in her own study.

Our estimate of the density of *M. siberu* is similar to that of Paciulli (2004) for *M. pagensis* in 10–20 year-old logged forests on North and South Pagai but is somewhat higher than those for undisturbed habitats. Nevertheless, we consider our estimate to be conservative, given the difficulties in obtaining reliable counts of group sizes in this species, in which groups can be relatively large and variable. It is therefore possible that we underestimated group size and that actual densities for the species may be >16 km⁻². Other data for Siberut itself are not available and comparisons with other macaque species in the Sundaland region are not useful because of the extreme degree of variation (<5 to >120 km⁻²) reported (MacKinnon, 1986).

Density estimates of *S. concolor*, *P. potenziani* and *H. klossii* are all higher in the Peleonan forest than in any of the locations surveyed on North and South Pagai by Paciulli (2004). The high primate densities we found in the Peleonan forest may reflect its relatively undisturbed structure and low hunting pressure by humans. Assuming that data from the main study site are representative of the 40 km² Peleonan forest as a whole (despite the inclusion of additional swamp forest habitat), approximate population sizes in the region could be 2,100 *S. concolor*, 600 *M. siberu*, 360 *H. klossii*, and 330 *P. potenziani*. However, these

estimates need to be verified by further surveys. Our data nevertheless shows that the Peleonan forest supports a substantial proportion of the global populations of all four Siberut primates and hence is of major importance for the conservation of Mentawai's endemic primates.

The threats posed to Mentawai's primates by logging activities have been documented for >20 years (Watanabe, 1981; Fuentes & Ray, 1995; Paciulli, 2004; Whittaker, 2005b). Whilst effects of forest destruction through logging are likely to be species-specific and recovery after logging can only be detected in time-lagged responses, all Mentawaiian primate species, and especially the two species of langur, appear to be negatively affected by logging at even low cutting intensities (Whitten, 1982b; Marsh *et al.*, 1987; Paciulli, 2004). In North Siberut logging and deforestation have so far not been as destructive as on North and South Pagai, and there is still hope that ecologically sound land-use strategies, and proper management of Siberut National Park, will play a much larger role on this island than on North or South Pagai. However, given the uniqueness of the so far largely unexplored fauna and flora of the island, more primate surveys and other biodiversity research are required. In particular, we advocate that wildlife studies on Siberut be carried out in both principal forest types, i.e. freshwater and peat swamp forest and dipterocarp forest. Most of the swamp forests are situated within production forests and will be subject to logging, whereas most of Siberut National Park is dipterocarp forest. The consequences of logging, particularly of swamp forests, for Siberut's endemic wildlife, are still largely unknown. Because the Siberut Conservation Project already operates in production forest, encompassing both forest types, Project activities offer future possibilities for research and for the development of conservation strategies on Siberut and other Mentawai islands.

Acknowledgements

We thank Conservation International, the Critical Ecosystem Partnership Fund, Margot Marsh Biodiversity Foundation, and the German Primate Centre for partly funding this survey. We particularly thank our field guides Pak Nauli and Pak Tarzan, as well as Muhamad Agil, Christos Astaras, Simone Riedelbauch, Andrea Herrmann and Juliane Geyer for very valuable assistance.

References

- BROCKELMAN, W.Y. & ALI, R. (1987) Methods of surveying and sampling forest primate populations. In *Primate Conservation in the Tropical Rain Forest* (eds C.W. Marsh & R.A. Mittermeier), pp. 23–62. Alan R. Liss, Inc., New York, USA.
- BROCKELMAN, W.Y., REICHARD, U., TREESUCON, U. & RAEMAEKERS, J.J. (1998) Dispersal, pair formation and social structure in gibbons (*Hylobates lar*). *Behavioural Ecology and Sociobiology*, 42, 329–339.

- BUCKLAND, S.T., ANDERSON, D.R., BURNHAM, K.P., LAAKE, J.L., BORCHERS, D.L. & THOMAS, L. (2001) *Introduction to Distance Sampling: Estimating Abundance of Biological Populations*. Oxford University Press, Oxford, UK.
- CHASEN, F.N. & KLOSS, C.B. (1927) Spolia Mentawiensia – Mammals. *Proceedings of the Zoological Society of London*, 53, 797–840.
- CHIVERS, D.J. (1986) South-east Asian Primates. In *Primates: The Road to Self-sustaining Populations* (ed. K. Benirschke), pp. 126–151. Springer, New York, USA.
- FUENTES, A. & RAY, E. (1995) Humans, habitat loss and hunting: the status of the Mentawai primates on Sipora and the Pagai islands. *Asian Primates*, 5. <http://pin.primate.wisc.edu/news/journals/Asian.5.3.html> [accessed 27 February 2008].
- IUCN (2007) *IUCN Red List of Threatened Species*. IUCN, Gland, Switzerland <http://www.iucnredlist.org> [accessed 5 February 2008].
- KITCHENER, A.C. & GROVES, C. (2002) New insights into the taxonomy of *Macaca pagensis* of the Mentawai islands, Sumatra. *Mammalia*, 66, 533–542.
- MACKINNON, K. (1986) The conservation status of nonhuman primates in Indonesia. In *Primates: The Road to Self-sustaining Populations* (ed. K. Benirschke), pp. 99–126. Springer, New York, USA.
- MARSH, C.W., JOHNS, A.D. & AYRES, J.M. (1987) Effects of habitat disturbance on rain forest primates. In *Primate Conservation in the Tropical Rain Forest* (eds C.W. Marsh & R.A. Mittermeier), pp. 83–107. Alan R. Liss, Inc., New York, USA.
- MCCONKEY, K.R. & CHIVERS, D.J. (2004) Low mammal and hornbill abundance in the forests of Barito Ulu, Central Kalimantan, Indonesia. *Oryx*, 38, 439–447.
- MITCHELL, A.H. & TILSON, R.L. (1986) Restoring the balance: traditional hunting and primate conservation in the Mentawai Islands, Indonesia. In *Primate Ecology and Conservation* (eds J.G. Else & P.C. Lee), pp. 249–260. Cambridge University Press, Cambridge, UK.
- MITTERMEIER, R.A., RATSIMBAZAFY, J., RYLANDS, A.B., WILLIAMSON, L., OATES, J.F., MBORA, D. *et al.* (compilers) (2007) *Primates in peril: the world's 25 most endangered primates 2006–2008*. *Primate Conservation*, 22, 1–40. Also available at <http://www.primate-sg.org/PDF/Primates.in.Peril.2006-2008.v2.pdf> [accessed 29 April 2008].
- NIJMAN, V. (2004) Conservation of the Javan Gibbon *Hylobates moloch*: population estimates, local extinctions and conservation priorities. *The Raffles Bulletin of Zoology*, 52, 271–280.
- PACIULLI, L.M. (2004) *The effects of logging, hunting and vegetation on the densities of the Pagai, Mentawai Island, primates*. PhD thesis, Stony Brook University, New York, USA.
- PERES, C.A. (1999) General guidelines for standardizing line-transect surveys of tropical forest primates. *Neotropical Primates*, 7, 11–16.
- PLUMPTRE, A.J. & COX, D. (2006) Counting primates for conservation. *Primates*, 47, 65–73.
- ROOS, C., ZIEGLER, T., HODGES, K.J., ZISCHLER, H. & ABEGB, C. (2003) Molecular phylogeny of Mentawai macaques: taxonomic and biogeographic implications. *Molecular Phylogenetics and Evolution*, 29, 139–150.
- ROWE, N. (1996) *The Pictorial Guide to the Living Primates*. Pogonias Press, New York, USA.
- SCHEFOLD, R. (1988) *Lia: Das große Ritual auf den Mentawai-Inseln (Indonesien)*. Reimer Verlag, Berlin, Germany.
- TENAZA, R.R. (1987) The status of primates and their habitats in the Pagai islands, Indonesia. *Primate Conservation*, 8, 104–110.
- TENAZA, R.R. & FUENTES, A. (1995) Monandrous social organization of pigtailed langurs (*Simias concolor*) in the Pagai Islands, Indonesia. *International Journal of Primatology*, 16, 295–310.
- TILSON, R. (1977) Social organization of simakobu monkeys (*Nasalis concolor*) in Siberut island, Indonesia. *Journal of Mammalogy*, 58, 202–212.
- THOMAS, L., LAAKE, J.L., STRINDBERG, S., MARQUES, F.F.C., BUCKLAND, S.T., BORCHERS, D.L. *et al.* (2004) *Distance 4.1 Release 2*. Research Unit for Wildlife Population Assessment, University of St. Andrews, UK. <http://www.ruwpa.st-and.ac.uk/distance/> [accessed 5 February 2008].
- VERSTAPPEN, H.T. (1973) A geomorphological reconnaissance of Sumatra and adjacent islands. Wolters-Noordhoff, Groningen, The Netherlands.
- WALTERT, M., LIEN, FABER, K. & MÜHLENBERG, M. (2002) Further declines of threatened primates in the Korup Project Area, south-west Cameroon. *Oryx*, 36, 257–265.
- WATANABE, K. (1981) Variations in group composition and population density of the two sympatric Mentawai leaf monkeys. *Primates*, 22, 145–160.
- WHITTAKER, D.J. (2005a) *Evolutionary genetics of Kloss's gibbons (Hylobates klossii): systematics, phylogeography, and conservation*. PhD thesis, The City University of New York, New York, USA.
- WHITTAKER, D.J. (2005b) New population estimates for the endemic Kloss's gibbon *Hylobates klossii* on the Mentawai islands, Indonesia. *Oryx*, 39, 458–461.
- WHITTEN, A.J. (1982a) Home range use by Kloss gibbon (*Hylobates klossii*) on Siberut Island, Indonesia. *Animal Behaviour*, 30, 182–198.
- WHITTEN, A.J. (1982b) *The Gibbons of Siberut*. Dent, London, UK.
- WWF (1982) *Siberut Nature Conservation Area, West Sumatra, Management Plan 1983–1988*. A Report for the Directorate of Nature Conservation, Bogor, Indonesia.
- YANUAR, A., FUENTES, A. & STUDD, K. (1998) A short report on the current status of the Mentawai snub-nosed pig tail langur (*Simias concolor concolor*) on Simalegu island, South Pagai, Mentawai, Indonesia. *Tropical Biodiversity*, 5, 99–305.

Biographical sketches

MATTHIAS WALTERT studies the effects of forest land use on tropical biodiversity and teaches wildlife population and biodiversity assessment within the integrated bi-national Msc *International Nature Conservation* at Georg-August-Universität Göttingen, Germany, and Lincoln University, New Zealand. CHRISTOPHE ABEGB is a specialist in macaque evolution and sociobiology. His work in Indonesia led to the instigation of the Siberut Conservation Project in 2002. THOMAS ZIEGLER's research interests include mating strategies and ecological adaptations of primates, primarily langurs. DODI PRIATA is the field manager of the Siberut Conservation Project. KEITH HODGES has research interests in evolutionary endocrinology and comparative reproduction in primates.